


The Lewistown Hospital Vascular Laboratory
 400 Highland Avenue, Lewistown, PA 17044
 Telephone 717-242-7031
 Philip L. Rice, M.D., F.A.C.S., R.P.V.I. Medical Director



Wells Pretest Probability for DVT

Clinical Feature

- Active cancer (treatment ongoing, within previous 6 months, or palliative) 1
- Paralysis, paresis, or recent plaster immobilization of the lower extremities 1
- Recently bedridden for > 3 days or major surgery within 4 weeks 1
- Localized tenderness along the distribution of the DEEP venous system 1
- Entire leg swollen 1
- Calf swelling > 3 cm when compared with the asymptomatic leg* 1
- Pitting edema (greater in the symptomatic leg) 1
- Collateral superficial veins (non-varicose) 1
- Alternative diagnoses as likely as or greater than that of DVT -2

Total

* Measured 10 cm below tibial tuberosity
 In patients with symptoms in both legs, the more symptomatic leg should be used.

High pretest probability	≥3 points
Moderate pretest probability	1-2 points
Low pretest probability	zero or negative points

Wells Pretest Probability for PE

Clinical Feature

- Suspected DVT 3.0
- An alternate diagnosis is less likely than PE 3.0
- Heart Rate > 100 beats/min 1.5
- Immobilization or surgery in the previous 4 wks 1.5
- Previous DVT/PE 1.5
- Hemoptysis 1.0
- Malignancy (on treatment, treated in past 6 mo, or palliative) 1.0

High pretest probability	>6 points
Moderate pretest probability	2-6 points
Low pretest probability	<2 points

Total
